

Report of the Coastal States Working Group
on the distribution of Norwegian spring
spawning herring in the North-East Atlantic
and the Barents Sea

October 2025

Table of Contents

1.	Introduction.....	2
1.1.	Terms of Reference.....	2
1.2.	Approach of the present group	2
2.	Background.....	3
2.1.	Main characteristics of the migration patterns of the stock.....	3
2.1.1.	Early life stages.....	3
2.1.2.	Adults.....	3
2.2.	Mixing with other herring stocks and populations	4
3.	Methods	5
3.1.	Zonal database	5
3.2.	Surveys	5
3.2.1.	Methods used when calculating abundance/biomass	5
4.	Results derived from surveys.....	6
4.1.	Adults from acoustic survey on the spawning grounds	7
4.2.	Larvae survey.....	7
4.3.	Juveniles in Barents Sea from the International ecosystem survey in the Nordic Seas (IESNS)	7
4.4.	Adults from the International ecosystem survey in the Nordic Seas (IESNS)	7
4.5.	0-group and juveniles from ecosystem survey in Barents Sea	8
4.6.	Adults from the International ecosystem summer survey in Nordic Sea (IESSNS)	8
5.	Results derived from catches	8
5.1.	Overview of submitted data.....	8
5.2.	Description of Fishery	9
5.3.	Seasonal and interannual patterns.....	9
6.	Discussion and conclusions	10
7.	References.....	12
8.	Tables.....	14
9.	Figures	22
	Annexes	56
	Annex 1 – Survey and catch tables from the 2014 report.....	56
	Annex 2. Distribution of NSSH fishing per month and year.....	56
	Annex 3. Data call and formats.....	69
	Annex 4. List of participants.....	71

1. Introduction

At the Coastal States meeting on 23-24 October 2019 between the European Union, the Faroe Islands, Iceland, Norway and the Russian Federation, the Norwegian delegation announced of its initiative where the Institute of Marine Research in Norway would lead a participatory process to update the "Report of the Coastal States Working Group on the distribution of Norwegian spring spawning herring in the North-East Atlantic and the Barents Sea" from 2014.

In the following Coastal states meeting in 2020 it was agreed that the report should be updated on an annual basis. The present report builds on the 2024 report with added information from catches in 2024 and surveys in 2024-2025.

In the agreed record from the 2021 consultations, the work of updating the report was formalized by establishing a working group. From the year 2023 onwards, the group should be chaired by a scientist from the party chairing the Coastal State Consultations, which is Faroes in 2025.

1.1. Terms of Reference

The text below is copied from the agreed record from the Coastal States consultations in 2020 (item 5) and is regarded as terms of reference for the present working group:

"The delegations welcomed a presentation from The Norwegian Institute of Marine Research on the progress with the participatory process to update the "Report of the Coastal states Working Group on the distribution of Norwegian Spring Spawning Herring in the North East Atlantic and the Barents Sea, Copenhagen 4-7 March 2014" and the overview of methods for calculations of zonal attachment. Furthermore, methodologies to be used for calculations of zonal attachment for this stock were presented. In this regard, the delegations welcomed the fact that Norway intends to continue this work and to convene other scientific meetings with a view to keep the report up-to-date."

1.2. Approach of the present group

The present report relies on standardised information in the form of output from nationally and internationally coordinated surveys with common databases, most of them annually reported to ICES.

For temporal distribution among zones the present report followed a direct observation approach like the 2014 WG. However, for the survey data after 2013, StoX (Johnsen et al., 2019) replaced the software BEAM that is no longer supported. The group was of the general opinion that the former analysis made using BEAM are comparable with StoX. The exercise to compare the few years of surveys data using both BEAM and StoX has only resulted in minor differences, therefore these can be considered as comparable.

Surveys provide snapshots of distribution of biomass/abundance of the different life stages. The former report used the following stages: early larvae, 0-group, juveniles (ages 0-3) and adults (age 4+), by producing maps and, when the data allowed, tables showing proportions by zones. The surveys provide information about biomass or abundance at specific points in time and space, but the integrated nature of their analysis will give distribution maps within the survey period. The present report contains added survey information from surveys in 2023-2025.

Catch distributions are based on a data call addressing each country with official recorded catches from the stock from logbooks and sales slips. This report contains added information from catches in 2024. The information from the landings statistics was compared to catches provided for ICES WGWIDE (ICES, 2025) and Coastal States. Data were requested and reported by year, month, country and zone as tonnes and proportions (Annex 3).

2. Background

2.1. Main characteristics of the migration patterns of the stock

The Norwegian spring-spawning herring (*Clupea harengus*) is one of the stocks composing the Atlanto-Scandian herring, together with two Icelandic herring stocks; the summer-spawners and spring-spawners (Johansen 1919; Dragesund et al., 1980). It is the largest herring stock in the world. It is widely distributed and highly migratory throughout large parts of the NE Atlantic during its lifespan. By far the majority of the adult stock occurs in Divisions 2a,b, 5a,b and 14a,b (Figure 2.1.1). Juveniles of the stock have their nurseries in Division 1a,b. In some years, Norwegian spring spawning herring have been found in adjacent areas mixing with other herring stocks.

The herring spawns along the Norwegian west coast in February-April. Large variations in the north-south distribution of the spawning areas have been observed through the centuries, and the most recent variation has been driven by the large 2016 year class that has shifted the spawning area northwards to the Lofoten area (Slotte et al., 2025). The larvae drift north and northeast and distribute as 0-group in fjords along the Norwegian coast and in the Barents Sea. The Barents Sea is by far the most important juvenile area for the large year-classes, which form the basis for the large production-potential of the stock. Some year-classes are in addition distributed into the Norwegian Sea basin as 0-group. Most of the young herring leave the Barents Sea as 3 years old and feed in the north-eastern Norwegian Sea for 1–2 years before recruiting to the spawning stock. With maturation, the young herring start joining the adult feeding migration in the Norwegian Sea. The feeding migration starts just after spawning with the maximum feeding intensity and condition increase occurring from late May to July (Homrum et al., 2022). The feeding migration is in general length dependent; the largest and oldest fish perform longer and typically more western migrations than the younger ones. Young fish of productive year-classes emerging from the Barents Sea can migrate to the northeast of the Norwegian Sea for feeding. After the dispersed feeding migration, the herring aggregates through September-October in one or more wintering areas. These areas are unstable and since 1950 the stock has used at least 6 different wintering areas in different periods. During the 1950s and 1960s they were situated east of Iceland and after around 1970 in Norwegian fjords. In 2001–2002 a new wintering area was established off the Norwegian coast between 69°30'N and 72°N in addition to the fjords. In 2007-2009, however, no herring was observed in the fjords in winter and a new wintering area was located in the ocean off northern Norway. After wintering, the spawning migration starts around mid-January. In the last almost two decades, the herring has partially stayed in the south-western part of the Norwegian Sea for a prolonged time in autumn.

2.1.1. Early life stages

Norwegian spring spawning herring spawns on the Norwegian coastal banks (Figure 2.1.1.1) during February-April (peak spawning in March). The eggs are deposited on the bottom. After hatching, larvae appear in the upper layers of the sea and drift with the current along the slope of the continental shelf. Larvae are gradually spread along the Norwegian coast and later appear in the Barents Sea along the Russian coast and sometimes the Bear Island-Spitsbergen area. While poor year-classes are mainly restricted to fjords in Norway, abundant ones are predominantly found in the Barents Sea. For instance, the portion brought to the Barents Sea of the abundant herring year-classes 1991, 1992 and 1998, comprised 84, 96 and 98 %, respectively, of the total number of these year-classes at the age of 1 year. Corresponding figures for the medium year-classes of 1990, 1993 and 1996 were 29-48 %, and of the poor year-classes 1995-1997 only 4-9 % (Krysov and Røttingen 2011). The young herring remain in the Barents Sea usually 3 years and then migrate westwards into the Norwegian Sea, but sometimes good year classes, as for example 2016, remain in the Barents Sea for 4-5 years.

Migration of the young herring in the Barents Sea varies seasonally. This has been described in detail for the 1983 year-class (Røttingen 1990) (Figure 2.1.1.2).

2.1.2. Adults

A characteristic feature of this herring stock is an extensive and varying migration pattern. The migration may be relatively stable for periods while periods of large changes occur occasionally at

varying time intervals. The changes may be observed on the spawning-, feeding- and/or overwintering grounds both in a single or an interrelated manner.

The usual annual clockwise migration route of adult herring feeding in the Norwegian Sea starting in the south after spawning and ending in the northeast at the wintering areas has persisted through the whole period. However, some changes in the distribution and recently the timing of the migration have been observed through this period.

After the recovery of the NSS herring stock, and until about 2002 the bulk of the adult herring wintered in fjords in northern Norway. The 1998 and 1999 year-classes were expected to enter the fjords around 2002 but were instead observed wintering off the coast in the ocean off Vesterålen/Troms, between 69°30'N and 72°N. This continued in the years to come and in 2005 also the 2002 year-class was observed wintering in the same area. During these years, the amount of older herring wintering in the fjords decreased rapidly and following the winter of 2007 no herring was observed in the fjords. In the last several years wintering herring have been observed in the oceanic areas and fjords in the Troms region.

The main spawning grounds of NSS herring are along the Norwegian coast from Siragrunnen in the south to Vesterålen in the north, with the main activity off Møre (Dragesund et al. 1980). Historically, the utilization of the different spawning grounds has been highly dynamic. In recent years the spawning has taken place further north than usual, such that since 2023 almost no herring spawned south of the Lofoten area (see Figure 4.1.1).

After spawning, in March/April the herring migrate westwards into the Norwegian Sea to start feeding and main concentrations are found in the central part of this area. In July, the herring are spread out over a wide area feeding around the fringes of the Norwegian Sea. In the early period from 1995-1999 the main concentrations of feeding herring utilised the central and southern Norwegian Sea, mainly the southern part of the International and the Norwegian zone. From 1999 a north-eastward displacement of the adult stock took place, which lasted until 2005, with almost no herring observed in the central region. In 2006-2016 they were more pronounced in the southern and south-western area. After 2016, the main concentration again shifted to the central and southern Norwegian Sea and the eastern part also. The last few years the southern boundary of the feeding distribution of herring in the Norwegian Sea has again shifted a bit further north to around 65 degrees north (see Figure 4.6.1, that shows the distribution of the feeding herring in July-August 2025).

It is not clear what drives the changes in the migration, but the biomass, age structure and production of zooplankton are likely factors, as well as oceanographic features such as sea temperature. A further factor might be the large pelagic stocks of mackerel and blue whiting occupying the same feeding area. Young herring leaving the nursery areas in Barents Sea or coastal waters in Norway and joining the adult stock for feeding is not as capable of extensive feeding migration as older herring. Correspondingly, when rich year-classes have joined the adult stock from the nursery areas in the Barents Sea a more north-easterly distribution has been observed. The stock is currently dominated by the 2016 year-class which was distributed over much of the Norwegian Sea. However, it is expected that the rich 2022 and 2021 year-classes will join the adult stock in the oceanic areas in 2026, and this might change the current distribution pattern of the herring.

2.2. Mixing with other herring stocks and populations

In the Norwegian Sea (2.a, 2.b), all monitored, sampled, and caught herring are assessed and managed as Norwegian spring-spawners. During the spawning season (1 February – 30 April), the distribution area is extended, and herring caught along the Norwegian coast (within the 12 nm) in divisions 4.a, can be recorded as Norwegian spring-spawners clearly split out from other populations in the North Sea based on maturation, age, length and vertebrae counts, in addition to the very specific spatiotemporal use of historic spawning grounds for NSS herring and evidence from tag-recapture data. During feeding migration, the Norwegian spring-spawners have the potential to mix with herring stocks and populations that have been identified based on spawning time and area, or otolith characteristics. In the Norwegian Sea and adjacent waters, they can mix with the Norwegian autumn-spawners (Husebø et al 2005), the Icelandic summer-spawners, the Icelandic spring-spawners (small component, Óskarsson 2018), the Faroese autumn-spawners (Figure 2.2.1), North Sea herring and components of the western Baltic herring stock (Bekkevold et al. 2023) and several local herring populations in Norway: in fjords along the coast (Aasen 1952; Lie et al 1978; Jørstad et al 1994; Libungan et al 2015), in semi-enclosed coastal ecosystems (Johannessen et al 2009; Langgård et al 2014) and in the brackish Lake Landvik in the south (Eggers et al 2014; Libungan et al 2015). In Iceland, the Norwegian spring-spawners and

Icelandic summer-spawners are separated based on maturity stage for management purposes, and in the Faroes they are separated from local autumn-spawners based on otolith structure and maturity stage, also for management purposes (i.e. splitting of catches).

3. Methods

3.1. Zonal database

In the analyses presented in this report, the same database with EEZ coordinates has been used as in the 2014 report (Anon. 2014). The countries provided their information on catches on NSS herring by EEZ, i.e. it was not necessary to use the database to split catches into EEZs. In the report from 2014 zonal information from the database was applied to survey data (Anon. 2014). However, the method used to produce the estimates of the surveys in the present report has changed. A new estimation software StoX (Johnsen et al., 2019) is now available where biomass estimates are produced directly by EEZs. For the records, please note that the shapefile (MarineRegions.org) used in StoX for the EEZs does not include disputed areas where more than one party claim certain rights. With regards to NSS herring this applies to a disputed area along the boundary between the Icelandic and the Faroese EEZs.

The StoX software is used as the estimation procedure in several ICES-coordinated surveys for the assessment of various stocks (e.g. the International Ecosystem Survey in the Nordic Seas – NSSH, the International blue whiting spawning stock survey – blue whiting, and the International Ecosystem Summer Survey in the Nordic Seas – NEA mackerel). This estimation procedure replaced the BEAM software, which produced biomass estimates on 2x2 squares of 1-degree latitude and 2 degree's longitude (instead of exact location). This new method is considered to provide more accurate estimation and thereby being more adequate than the former one. A comparison between the two methods was made on the estimates for two years for all the international surveys and it showed minor differences in most cases. Re-estimating the whole time series back to 1995 is not currently possible because some of the acoustic data are not available in survey databases/repositories (PGNAPES / ICES acoustic database) prior to 2009.

3.2. Surveys

3.2.1. Methods used when calculating abundance/biomass

Larval index

The herring larval abundance index is based on the number of caught larvae per square metre of surface at each station and is calculated with consideration to measured water volume, depth layers and total number of caught larvae. All larvae are included in the index and herring larvae abundance indices are produced. The surveyed area is delimited by a polygon, which is divided into a grid with cell sizes of 14.5*8 km. The herring larval density per grid is estimated and summed over the covered area (Stenevik *et al.* 2012).

0-group index

The geographical distribution of 0-group fishes is estimated by the standard procedure which was first recommended in 1980 (Anon. 1980). All vessels use a small mesh mid-water trawl (“Harstadtrål”). The standard procedure consisted of tows at 3 depths, each of 0.5 nautical miles, with the headline of the trawl located at 0, 20 and 40 m. When the 0-group fish layer was recorded on the echo-sounder deeper than 60m or 80m additional tows at 60 and 80m, of 0.5 NM distance also, were carried out. The history of development of 0-group investigation and assessment method is described in detail for example in the survey report from 2007 (e.g. Anon. 2007). The abundance of 0-group herring is measured in numbers and the index is calculated using the StoX software.

Juveniles

Juveniles are defined as immature herring age 1-3. All regions of the Barents Sea and adjacent areas of the Norwegian Sea are covered, with transects 35 nautical miles apart. Data from pelagic trawl hauls and bottom trawl hauls considered representative for the pelagic component of the stocks, which is

measured acoustically, are included in the stock abundance calculations. The StoX software is used to make estimates of total biomass and numbers of individuals by age and length in the whole survey area and within different subareas.

Adults

The survey estimates of the adult part of the stock are based on scientific echosounders combined with trawl catches to identify species and size distribution. The StoX software is used to make estimates of total biomass and numbers of individuals by age and length in the whole survey area and within different subareas.

4. Results derived from surveys

A number of national and international research surveys have been conducted over the period 2014-2025 that can provide information about spatial and temporal variation in distribution and quantity of different life stages of Norwegian spring-spawning herring (Table 4.1). Some of the surveys are constrained to only part of the distribution area, while others provide information for the whole stock and/or the whole period from 2014 to 2024/2025. All surveys are snap shots and representative for the distribution during the survey period. Seven survey series are updated since the 2014 WG (only six from 2017) and these surveys, taking place in the first to third quarters of the year, are considered to provide adequate quantitative information about the distribution of the stock, adult or juveniles. Relative estimates on biomass or number of fish for different EEZs were tabulated. No survey information is available for quarter 4. Below is a short description of each of the surveys, their main results with regards to inter-annual variation in the spatial distribution of the stock, and conclusions that can be derived from them as well as their limitations.

Table 4.1. List of relevant surveys targeting Norwegian spring spawning herring. Indication is given of whether and how the measured abundance/biomass is representative for the distribution of the life-stage. Some surveys may have started earlier, but the first year under consideration in this report is 2014 (the former reports include surveys in the period 1995-2013, e.g. Anon. 2024).

Survey	Month	Life stage	Complete spatial coverage of life-stage	Year range	Representative for distribution of life-stage
1st quarter					
Acoustic survey on the spawning grounds	Feb/ Mar	Adults	Yes	2015-2025	Whole spawning stock covered
2nd quarter					
International ecosystem survey in the Nordic Seas (IESNS) – Barents Sea	May	Juveniles ages 1-3	Yes	2014-2024	Main distribution area of juveniles covered
International ecosystem survey in the Nordic Seas (IESNS)	May	Adults	Yes	2014-2025	Adult part of stock spatially covered
3rd quarter					
Ecosystem survey in Barents Sea	Aug/O ct	0-group,	Yes	2014-2024	Main area of 0-group covered
	Aug/O ct	Juveniles age 1-3	Yes	2014-2024	Main area of juveniles covered
International Ecosystem Summer Survey in the Nordic Seas (IESSNS)	Jul/ Aug	Adults	Yes	2014-2025	Northern boundary of distribution area not fully reached in all years Main distribution area covered
4th quarter					
					No surveys

1st quarter

4.1. Adults from acoustic survey on the spawning grounds

A Norwegian acoustic survey has been undertaken to estimate the abundance at age of herring in the spawning areas along the Norwegian coast in February and March. The survey has been carried out since 1988 but not in every year. This survey is conducted during a period when the entire spawning stock is distributed along the Norwegian coast to spawn (Figure 4.1.1). Consequently, 100% of the spawning stock is within Norwegian EEZ in this period. No surveys were conducted during 2009 – 2014. In 2015 the survey was re-introduced, and results from 2015 – 2025 are included in this report.

2nd quarter

4.2. Larvae survey

A Norwegian herring larvae survey was carried out on the Norwegian shelf 1981 – 2016 during March-April. The objectives of the survey were to map the distribution of herring larvae and other fish larvae on the Norwegian shelf and to collect data on hydrography, nutrients, chlorophyll and zooplankton (Figure 4.2.1). There have been no surveys after 2016.

4.3. Juveniles in Barents Sea from the International ecosystem survey in the Nordic Seas (IESNS)

A part of the international ecosystem survey in the Nordic Seas in May (section 4.6), is focusing on juvenile herring, zooplankton and hydrography in the Barents Sea. The survey is conducted by a Russian vessel and has been carried out in most years since 1995. There was no survey in 2020, 2022, 2023 or 2025. Results from 2014 – 2024 are included in this report. The distribution maps are presented in Figure 4.3.1 and percentages by zone in Table 4.3.1. Average percentage distribution (2014 – 2021 and 2024) is: Russia 58.2%, Norway 41.8%.

4.4. Adults from the International ecosystem survey in the Nordic Seas (IESNS)

The international ecosystem survey in the Nordic Seas is aimed at observing the pelagic ecosystem, focusing on herring, blue whiting, zooplankton and hydrography. The survey, carried out in late April and May since 1995, is coordinated by ICES survey planning groups (PGNAPES and currently WGIPS) and is a cooperative effort by Faroes, Iceland, Norway, Russia, and since 1998 the EU (Denmark, Germany, Ireland, The Netherlands, Sweden and UK). Since 2022 the UK has also participated. A good internal consistency among years for age groups 4+ and expert judgment on the whole time-series point out that the survey covers the adult part of the stock adequately each year. Biomass estimates based on the acoustic and trawl samples for the years 2014-2025, shown in Figure 4.4.1 and Table 4.4.1, are therefore considered to be representative for the relative distribution of the adult part of the stock during that month of the year.

The average percentage distribution by zones for the whole period 2014 – 2025 give the following (Table 4.4.1): Norway 29.9%, International zone 20.2%, Faroes 19.5%, Iceland 23.3%, Jan Mayen 4.4%, and EU/UK 2.7%.

3rd quarter

4.5. 0-group and juveniles from ecosystem survey in Barents Sea

The survey consists of a trawl survey catching 0–group herring amongst other species and an acoustic survey estimating 1–3-year-old herring. It is difficult to assess the abundance during autumn, for various reasons. The age groups 1 to 3 are found mixed with 0–group herring and are difficult to catch in the sampling trawl used in this survey. The stock size estimates of herring are therefore considered less reliable than those for capelin and polar cod. The survey, however, is assumed to cover the main distribution area of 0–group herring even though it is known that they are found to a lesser extent in Norwegian fjords. In some years there is no coverage in the Russian EEZ. Distribution maps of the 0–group are presented in Figure 4.5.1. The results from this survey, tabulated for the years 2014–2024 for 0–group, shows an average distribution of 49.9% in Norwegian EEZ and 3.4% in Russian EEZ, 39.0% in the Fishery protection zone around Svalbard, and 8.6% in international waters (Table 4.5.1).

For the juveniles, the survey coverage was considered adequate in 2024, and results are presented in Figure 4.5.2. According to Table 4.5.2 an average distribution of the juveniles is 59.8% in Norwegian EEZ, 37.3% in Russian EEZ.

4.6. Adults from the International ecosystem summer survey in Nordic Sea (IESSNS)

This ecosystem survey was initiated in 2004 by Norway and has since been gradually expanded in geographical coverage, especially from 2009 and onwards with participation of vessels from Iceland and the Faroes in addition to two vessels from Norway and in the most recent years Greenlandic and Danish vessels. The main objective of the survey is to study abundance and distribution of Northeast Atlantic mackerel, NSSH, blue whiting and other pelagic species with acoustic and swept-area methods in relation to oceanographic conditions, prey communities and marine mammals. Acoustic estimates of herring are available from the 2014 – 2025 surveys (Figure 4.6.1 and Table 4.6.1). According to Table 4.6.1 the biomass, on average, during July/August (2014–2025) is distributed in the EEZ of Iceland 34.2%, Faroe Island 18.8%, Norway 20.1%, Jan Mayen 10.3%, the International waters in the Norwegian Sea 8.9%, Svalbard 5.8%, EU/UK 1.7% and Greenland 0.2%.

4th quarter

No new survey information is available for 4th quarter.

5. Results derived from catches

5.1. Overview of submitted data

To update the distribution maps of the catches for 2024, a request was formulated on catch data by year, month and ICES statistical square (0.5° latitude, 1° longitude) and by economic zone (EEZ) to the nations fishing for Norwegian spring-spawning herring. The deadline for submission of the catch data was set to 1st June 2025. All countries/parties delivered their catch data in time to be included in the report. The level of detail of the catch data was in line with the request.

The total catch in the years 1995–2024 and the relative catch by zones is shown in Figure 5.1.1. In Table 5.1.1 the catches reported to the Working Group are compared with the catches reported to WGWIDE (ICES 2025) and to catches delivered by Coastal States for each year in 2013–2024. Catches reported to the Working Group as percentage of catches reported to WGWIDE by year varies from 97.4 % to 101.7 % (Table 5.1.1). Obviously, the difference can partly be explained with the fact that the catch data were raised in the WGWIDE report to account for changes in the percentages of water

content. The proportion is 99.61 % for the whole period. There are however differences in catches reported by Coastal States with other data sources.

All catch data submitted had the requested resolution, i.e. catches reported by year, month, zone and ICES rectangle.

5.2. Description of Fishery

The fishery today is carried out mainly in the beginning of the year and in the autumn. It is mainly by large purse seiners and pelagic trawlers, but also to a smaller degree as a coastal fishery. The catches are used for human consumption and reduction purposes to fish meal and fish oil.

The fishery is regulated and carried out by the Coastal States. The TAC is set by the Coastal States and derived from an agreed long-term management plan (ICES 2018). Since 2014, however, the Coastal States have not agreed on a comprehensive sharing arrangement with parties allocating autonomous quotas resulting in catches exceeding the advised TAC.

In 2013-2024 the fishing pattern corresponds to the general seasonal distribution of the herring in the Norwegian Sea as the year progresses. The fishery focuses on wintering, pre-spawning, spawning and feeding fish. The fishing activity starts in January on the Norwegian shelf. The fishery in spring and early summer is nearly non-existing. The fishing effort then shifts south-west to Icelandic and Faroese waters in early autumn and expands north-east to International waters and Norwegian zone in late autumn.

In the earlier period until 2010, the traditional fishing activities followed the stock in a north-eastern direction in autumn to the eastern part of the Norwegian Sea. However, in the last decade the fishing activity have been significant also in the southern Norwegian Sea, in international (<68°N), Icelandic and Faroese EEZs in this time of the year. Fishing activities by zones in 2013-2024 is shown in Table 5.2.1 and Table 5.2.2.

The NSSH changed wintering areas from the fjords to more offshore waters off northern Norway during the years 2002-2006. The change in wintering pattern caused a large change in fishing pattern as well. More catches were taken during the spawning migration and at spawning time instead of during the wintering period. These changes applied mostly to the Norwegian fleet. In recent years, part of the herring has been wintering in the fjords of northern Norway again.

In recent years the herring have spawned further north than previously with almost no spawning south of Lofoten area. This has also affected the fishery along the Norwegian coast and to a certain degree also the high seas fishery in the south with less catches in the southern areas.

Typically, catches in the purse seine and pelagic trawl fishery consist of only herring with limited by-catches. However, due to the changes in the distribution of mackerel in the late 2000s, by-catches of mackerel did increasingly occur on the traditional herring fishing grounds but seem to have decreased again the last decade.

Due to limitations, which are inherent in catch data, e.g. regulation effects, national agreements on zonal access, distance from homeport, area misreporting, technical changes, changes in fishing methods etc., the fisheries do not necessarily depict the distribution of herring in the Norwegian Sea. However, fisheries do show presence of herring while no fishery cannot per se be concluded as absence of herring.

5.3. Seasonal and interannual patterns

Changes have been observed in the fishery through the years since the NSS herring reappeared in the feeding areas in the Norwegian Sea in the mid-1990s. The limitations mentioned in section 5.2 also apply to the seasonal changes in the herring fisheries and therefore the following description. The following description is therefore restricted to general remarks on distribution of the fisheries.

The changes observed in the seasonal pattern since the fishery began in the mid-1990s is depicted in Figure 5.3.1 showing the percentage distribution of the fishery by month for each year since 1995, and in Annex 2 where maps of the fishery per month and per year is shown. From this, four main periods

stand out in the seasonal fishing patterns: an early period 1995-1997, a mid-period from 1998-2005, a late period (2006-2012) and a recent period (2013-2024, Figure 5.3.1).

In the early period, up to 1998, the main fisheries were in the spawning area (Norwegian EEZ) in January-March with a peak in February, then a fishery in the Norwegian Sea (ICES Divisions 5.a, 5.b and 2.a) developed with a peak in May, followed by a fishery in the north-eastern areas from September to November (Figure 5.3.1).

In the mid period 1998-2005 the spawning fishery was the same, but the peak in the summer fishery changed from May to June while the autumn fishery continued as usual (Figure 5.3.1). However, the effort shifted with less effort in the summer period to an increase in the autumn fishery. A further change during this period was that the summer fishery had shifted further north into the Jan Mayen, International and Norwegian areas (Figure Annex 2). This northern movement of the feeding stock during the 1998-2005 could be driven by a combination of increased temperature and reduced zooplankton abundance in the Atlantic water masses further south (Eliassen et al., 2021).

In the period after 2005 the summer feeding areas again shifted southwest as was the case in the early period. The fishery, however, did not follow the previous pattern as the May and June fishery has almost disappeared, instead the late autumn and winter fishery dominated (Figure 5.3.1).

The recent fishing pattern is now concentrated in January targeting pre-spawning fish with practically no catches in March. The summer fishery has ceased and the only nations fishing in late summer are the Icelandic and Faroese fleets. In the total picture these quantities are small. The level of the autumn fishery has increased in recent years with a peak in October - November amounting to around 60% of the total annual catches (Figure 5.3.1). This fishery has for some years been mainly in the central Norwegian Sea, north of the Faroes and east of Iceland, whereas before 2015 it used to be stretched out towards the coast of Norway and north towards the Bear Island. Since 2019, however, there was also a fishery on the 2016 year-class off northern Norway during autumn. Changes in migration have also resulted in late arrival at the Norwegian coast for part of the stock (mostly older fish) during the winter in recent years.

There are several possible explanations for these observed changes in the summer fishery for Norwegian spring spawning herring. The distribution of herring has changed throughout the summer months over the years, and thus e.g. accessibility to the fleets may have also changed. The quality and fat content increases throughout the summer season, which may delay the fishery to increase profit. Stock size and consequently quota sizes may also affect where and when the catches are taken. Finally, the quota and fishing opportunities for other pelagic species (blue whiting and mackerel) are also likely to influence the focus put on the herring fishery.

The usual clockwise migration route of adult herring feeding in the Norwegian Sea, starting in the south after spawning, migrating westwards through the Faroese area and moving further north-westwards into Icelandic area in summer, and further north into the Jan Mayen area later in the summer, and ending in the northeast at the wintering areas seem to have stopped in the last decade. Now part of the adult herring seems to prolong their feeding period into the autumn months in the southern and southwestern areas, and the return migration to the wintering areas off northern Norway seems to start in the southern/central Norwegian Sea rather late in the season (October/November) or even later in the year. However, as a fishery was going on in the northern area during autumn at the same time as in the southern area, this indicates that it is not the entire stock that delays the northward migration. This pattern can be seen in the fishery by month from 1995-2024 (Figure Annex 2).

6. Discussion and conclusions

In this report data are presented from 2013-2025, which illustrate the distribution, and changes in distribution, of NSS herring. Tables from the period 1995-2013 used to be included in the present report for comparison as Annex 1. These are now removed from the present report, and the reader is advised to refer to any previous report up to and including 2024 instead (Anon. 2024). The main source of information is surveys. The surveys provide information about the general seasonal migration pattern and distribution of NSSH. Survey information is not available for all months, and some of the survey-series have been terminated. For example, there are no surveys covering the adult part of the population in the fourth quarter of the year during the period 2014-2025. The lack of systematic

information throughout the year prevents a full assessment of the zonal distribution for all life stages, without making assumptions based on other data than survey data. For adults, survey information is available for February and the summer months (May and July-August), but no survey information was available to the working group for April, when the herring start the feeding migration, and September – December, when the herring is finishing the feeding season and returning to the wintering areas. For larvae and 0-group herring survey information is available in April and August/September, and for juveniles in May and August/September.

There are limitations in using catch-data to describe the seasonal distribution of herring. The most obvious one is that the catch data show presence of herring while absence needs to be based on judgement. Other limitations include fishery regulation effects, changes in fleet behaviour, national agreements on zonal access and area misreporting. One example of regulation effects is, that there has been no directed fishery in the Barents Sea since 1987 to protect juveniles. However, taking these limitations into account the catch data can be used to some extent to describe the distribution of NSS herring, as a supplement to survey data.

By combining the information from the surveys and the catches, some general conclusions can be made for the adult part of the stock and partly for juveniles, but a full assessment of a zonal distribution is not possible.

Information about the distribution of larvae, 0-group and juvenile stem from surveys only.

Larvae are distributed along the Norwegian coast in spring and drift mainly into the Barents Sea. That is, they are within the EEZ of Norway (larvae survey until 2016).

0-group herring were distributed in the Barents Sea in August in 2014-2025. There was some variability between years, but most being within the Norwegian zone. The rest is, in most years, distributed in the Russian zone, but also in the Fishery protection zone around Svalbard.

Juveniles were primarily distributed in the Barents Sea. In the years 2014-2025 information about the distribution in May comes from the IESNS surveys (the area was not surveyed in 2020, 2022, 2023 and 2025) and only the eastern part of the Barents Sea was covered in 2024. According to these surveys most of the biomass was within the EEZs of Russia and Norway. Joint Russian and Norwegian surveys have taken place in August/September in the years 2014-2024 and also in these surveys most of the juveniles were within the EEZs of Russia and Norway.

For the adults in quarter 1, the main sources of information are from surveys on the spawning grounds in 2015-2025. According to this information the distribution of the adult part of the stock is limited to the eastern part of the Norwegian Sea and the Norwegian fjords that is predominately within the Norwegian EEZ. Since around 2021, the spawning has been taken place progressively more at the northern spawning grounds with almost no spawning in 2025 on the main grounds in the south. In 2018, 2021 and 2022, catches indicated some herring in the International waters in quarter 1 (in January and February). No survey was conducted in 2014.

In quarter 2, the adult stock migrates from the spawning grounds to the feeding grounds. The IESNS survey in the years 2014-2025 shows some inter-annual variability in the distribution but no apparent trend. The biomass has been mostly distributed in four zones, namely, EEZs of Norway, Iceland, Faroes and International waters in the Norwegian Sea. The large 2016 year class had a more northeasterly distribution than older herring when recruiting to the spawning stock (~2019-2021) but has shown a similar pattern since then, although the distribution of the whole stock has been more northerly in the west the last three years. In 2025 young herring (2021 and 2022 year-classes) were present in the northeastern corner of the survey area.

Quarter 3 covers the latter half of the feeding season and the period when the stock starts to migrate back to the overwintering grounds. The results from the IESSNS survey in 2014-2025, are considered to be representative for the relative distribution of the adult part of the stock during July/August. The survey shows that herring is distributed mostly in the southwestern part of the Norwegian Sea in July/August, predominately in the EEZs of Iceland, Faroes and Norway. However in 2021 and 2022 a significant proportion was observed in the Jan Mayen area and further north into Svalbard waters in 2023. In 2025 highest proportions were in the northeastern corner of the survey area.

In quarter 4 there is no survey information for the period 2014-2024. However, data from the fishery indicate that the herring is widely distributed in several EEZs, at least during October and November. Moreover, the fisheries in the traditional feeding areas in the south-western part of the Norwegian Sea in quarters 3 and 4 have extended further into the autumn months, indicating that herring has been staying longer in this region compared to previous years. The reasons for the prolonged stay in the south-western regions of the feeding area in recent years are most likely linked to extended feeding opportunities into the autumn months; since around 2005 herring put on a large proportion of the weight gain after July and the autumn body condition is generally good (Homrum et al., 2022). However, in the last few years it seems as much of the adult herring stock (mainly the 2016 year-class) has migrated further north than the previous year-classes did during the oceanic phase.

The working group has collated available data from surveys in the period 2014-2025 and analysed distribution of NSS herring based on these surveys. Catch data from 2013 to 2024 have been collated as well and they supplement the survey data. The working group believes that the current report gives a relevant picture of the distribution – and changes in distribution – of NSS herring in the Northeast Atlantic in the period considered.

7. References

Aasen. O. 1952. The Lusterfjord herring and its environment. Reports on Norwegian Fishery and Marine Investigations 2, 1–63.

Anon. 1980. Preliminary report of the International 0-group fish survey in the Barents Sea and adjacent waters in August/September 1978. *Annals biol.*, Copenhagen, 35: 237-280.

Anon. 2007. Survey report from the joint Norwegian/Russian ecosystem survey in the Barents Sea in August-September 2007 (vol.1). IMR/PINRO Joint Report Series, No.4/2007.ISSN 1502-8828. 97 pp.

Anon. 2014. Report of the Coastal States Working Group on the distribution of Norwegian spring spawning herring in the North-East Atlantic and the Barents Sea. Copenhagen, 4 – 7 March 2014. 218 pp.

Anon. 2024. Report of the Coastal States Working Group on the distribution of Norwegian spring spawning herring in the North-East Atlantic and the Barents Sea. October 2024. 82 pp. https://assets.publishing.service.gov.uk/media/672b862940f7da695c921c4e/Coastal_States_NSSH_Distri_Report2024.pdf

Bekkevold, D., Berg, F., Polte, P., Bartolino, V., Ojaveer, H., Mosegaard, H., Farrell, E.D., Fedotova, J., Hemmer-Hansen, J., Huwer, B., Trijoulet, V., Albertsen, C.M., Fuentes-Pardo, A.P., Gröhsler, T., Pettersson, M., Jansen, T., Folkvord, A., Andersson, L. 2023. Mixed-stock analysis of Atlantic herring (*Clupea harengus*): a tool for identifying management units and complex migration dynamics, *ICES Journal of Marine Science*, 80, 173–184, <https://doi.org/10.1093/icesjms/fsac223>

Dragesund, O., Hamre, J., Ulltang, Ø. 1980. Biology and population dynamics of the Norwegian spring-spawning herring. *Rapp P-v Réun Cons int Explor Mer* 177 43-71.

Eggers, F., Slotte, A., Libungan, L.A., Johannessen, A., Kvamme, C., Moland, E., Olsen, E.M., Nash, R.D.M. 2014. Seasonal Dynamics of Atlantic Herring (*Clupea harengus* L.) Populations Spawning in the Vicinity of Marginal Habitats. *PLoS ONE* 9(11): e111985, doi 10.1371/journal.pone.0111985.

Eliassen, S.K., Homrum E.í., Jacobsen, J.A., Kristiansen, I., Óskarsson, G.J., Salthaug, A., Stenevik, E.K. 2021. Spatial Distribution of Different Age Groups of Herring in Norwegian Sea, May 1996–2020. *Frontiers in Marine Science*, 8: 1-13. doi: 10.3389/fmars.2021.778725

Homrum E. í, Óskarsson G. J., Ono K., Hølleland S. and Slotte A. 2022. Changes towards stable good somatic condition and increased gonad investment of Norwegian spring-spawning herring (*Clupea*

harengus) after 2005 are linked to extended feeding period. *Front. Mar. Sci.* 9:803171. doi: 10.3389/fmars.2022.803171

Husebø, A., Slotte, A., Clausen, L.A.W., and Mosegaard, H. 2005. Mixing of populations or year class twinning in Norwegian spring spawning herring? *Marine and Freshwater Research* 56: 763-772.

ICES. 2018. Report of the Workshop on a long-term management strategy for Norwegian Spring-spawning herring (WKNSSHMSE), 26–27 August 2018, Torshavn, Faroe Islands. ICES CM 2018/ACOM:53. 113 pp. <https://doi.org/10.17895/ices.pub.5583>.

ICES. 2025. Working Group on Widely Distributed Stocks (WGWIDE). ICES Scientific Reports. *In press*.

Johannessen, A., Nottestad, L., Ferno, A., Långard, L., and Skaret, G. 2009. Two components of Northeast Atlantic herring within the same school during spawning: support for the existence of a metapopulation? *ICES Journal of Marine Science*, 66: 1740-1748.

Johansen A.C. 1919. On the large spring-spawning sea herring (*Clupea harengus* L.) in the north-west European waters // *Medd. Komm. Havundersog., Ser. Fisk.*, 5 (8). 19 p.

Johnsen, E., Totland, A., Skålevik, Å., Holmin, A. J., Dingsør, G. E., Fuglebakk, E., & Handegard, N. O. 2019. StoX: An open source software for marine survey analyses. *Methods in Ecology and Evolution*. 10 :1523 –1528. <https://doi.org/10.1111/2041-210X.13250>

Jørstad, K.E., Dahle, C. and Paulsen, O.I. 1994. Genetic comparison between Pacific herring (*Clupea pallasi*) and a Norwegian fjord stock of Atlantic herring (*Clupea harengus*). *Canadian Journal of Fisheries and Aquatic Sciences* 51: 233-239.

Krysov, A.I., Røttingen, I. 2011. The Barents Sea: Ecosystem, Resources, Management. Half a century of Russian-Norwegian cooperation / IMR, PINRO. - Tapir Academic Press, Trondheim. - Chap. 5.3, 215-225.

Langgård, L., Fatnes, O.A., Johannessen, A., Skaret, G., Axelsen, B.E., et al. 2014. State-dependent spatial and intra-school dynamics in pre-spawning herring *Clupea harengus* in a semi-enclosed ecosystem. *Marine Ecology Progress Series* 501: 251-263.

Libungan, L., Slotte, A., Husebø, Å., Godiksen, J.A. and Pálsson, S. 2015. Latitudinal gradient in otolith shape among local populations of Atlantic herring (*Clupea harengus* L.) in Norway. *PloS one* 10; 6; e0130847. <https://doi.org/10.1371/journal.pone.0130847>

Lie, U., Dahl, O., and Østvedt, O.J. 1978. Aspects of the life history of the local herring stock in Lindåspollene, western Norway. *Reports on Norwegian Fishery and Marine Investigations*: 369–404.

Óskarsson, G.J. 2018. The existence and population connectivity of Icelandic spring-spawning herring over a 50-year collapse period. *ICES Journal of Marine Science*, 75: 2025–2032.

Røttingen I. 1990. The 1983 year class of Norwegian spring spawning herring as juveniles and recruit spawners//Biology and fisheries of the Norwegian spring spawning herring and blue whiting in the Northeast Atlantic//Proceedings of 4th Sov.-Norw. symp.– Bergen, Norway, 165-204.

Slotte, A., Salthaug, A., Vatnehol, S., Johnsen, E., Mousing, E. A., Høines, Å., Thorsen, C. B., Bjarnason, S., Homrum, E., Skagseth, Ø. and Stenevik E. K. 2025. Herring spawned poleward following fishery-induced collective memory loss. *Nature*, 642, pp 965–972. <https://doi.org/10.1038/s41586-025-08983-3>

Stenevik, E.K., Nash, R.D.M., Vikebø, F, Fossum, P. and Bakkeplass, K. 2012. The effects of survey design and circulation pattern on the perceived abundance of herring larvae: A case study for Norwegian spring spawning herring (*Clupea harengus*). *Fisheries Oceanography*, 21(5): 363-373.

8. Tables

Table 4.3.1. Percentages of NSSH by EEZ based on juveniles in Barents Sea during the International ecosystem survey in the Nordic Seas (IESNS) in May/June. No surveys conducted in 2020, 2022, 2023 and 2025. The coverage in 2024 was reduced and only the eastern part of the Barents Sea was covered.

Year	cruise	Norway	Russia	Svalbard	Total
2014		38.8	61.2	0.0	100.0
2015		51.1	48.9	0.0	100.0
2016		93.1	6.9	0.0	100.0
2017		26.3	73.7	0.0	100.0
2018		32.7	67.3	0.0	100.0
2019		30.8	69.2	0.0	100.0
2020					
2021		60.5	39.5	0.0	100.0
2022					
2023					
2024*		1.3	98.7	0.0	100.0
2025					
Average		41.8	58.2	0.0	

*Minimal coverage of the Norwegian EEZ

Table 4.4.1. Percentages of NSSH by EEZ based on adults during the International ecosystem survey in the Nordic Seas (IESNS) in May/June. *Change from EU to UK EEZ from 2021.

Coastal States WG Herring 2025

Year	EU/UK	Norway	Iceland	Svalbard	JanMayen	Faroes	InterNorwSea	Total
2014	2.9	21.8	14.0	0.0	0.8	31.0	29.6	100.0
2015	3.0	43.8	15.8	0.0	0.7	17.8	19.0	100.0
2016	9.5	23.7	16.7	0.0	3.6	27.3	19.2	100.0
2017	0.2	25.4	42.7	0.0	2.5	17.7	11.5	100.0
2018	2.3	14.3	30.3	0.0	5.8	28.5	18.8	100.0
2019	2.1	23.0	24.7	0.0	2.7	32.6	15.0	100.0
2020	1.5	43.9	21.4	0.0	0.1	27.8	5.2	100.0
2021	3.3	29.0	27.7	0.0	0.3	27.7	12.1	100.0
2022	1.0	40.2	20.4	0.0	5.1	9.4	23.9	100.0
2023	1.2	2.9	29.2	0.0	26.7	6.0	34.0	100.0
2024	2.1	45.2	16.3	0.0	1.3	5.7	29.3	100.0
2025	3.5	45.9	20.1	0.0	2.9	2.7	25.0	100.0
Average	2.7	29.9	23.3	0.0	4.4	19.5	20.2	

Table 4.6.1. Percentages of NSSH by EEZ based on adults during the International ecosystem summer survey in the Nordic Seas (IESSNS) in July/August. *Change from EU to UK EEZ from 2021.

Year	EU/UK	Norway	Iceland	Svalbard	JanMayen	Greenland	Faeroes	terNorwSe	Total
2014	2.4	3.7	61.8	0.1	8.3	0.0	23.0	0.8	100.0
2015	0.6	13.4	29.7	0.0	4.7	0.0	47.8	3.7	100.0
2016	0.1	5.6	66.5	0.0	1.6	0.3	25.6	0.2	100.0
2017	0.4	24.3	43.7	0.0	0.5	1.1	29.8	0.3	100.0
2018	6.6	20.6	52.7	0.7	0.4	0.0	18.7	0.3	100.0
2019	5.3	18.4	39.9	6.2	0.3	0.0	24.8	5.2	100.0
2020	0.2	26.8	27.5	19.3	2.2	0.0	15.2	8.8	100.0
2021	3.0	34.3	15.4	3.5	23.4	0.1	2.7	17.7	100.0
2022	1.2	14.6	14.5	9.2	18.5	0.4	20.6	21.0	100.0
2023	1.1	20.4	24.1	27.8	6.9	0.2	3.9	15.6	100.0
2024	0.0	9.5	14.6	1.2	42.9	0.0	6.9	25.0	100.0
2025	0.0	49.7	20.7	1.2	13.9	0.0	6.3	8.3	100.0
Average	1.7	20.1	34.2	5.8	10.3	0.2	18.8	8.9	

Table 5.1.1. Norwegian spring spawning herring. Comparison (%) of catches reported to the Coastal States zonal Working Group (WG data) and ICES WG WIDE (ICES. 2025). Coastal States catches are also shown (not available yet for 2024), which represent official catches, whereas WG WIDE catches are provided by scientists.

Year	WG data	WG WIDE	Comparison	Coastal States data
2013	684,776	684,743	100.0%	687,732
2014	449,199	461,306	97.4%	445,404
2015	320,128	328,740	97.4%	324,995
2016	384,697	383,174	100.4%	384,492
2017	720,081	721,566	99.8%	719,104
2018	594,367	592,899	100.2%	591,401
2019	773,565	777,165	99.5%	774,491
2020	720,796	720,937	100.0%	719,100
2021	848,099	851,813	99.6%	848,836
2022	827,686	813,834	101.7%	835,848
2023	678,583	680,552	99.7%	680,507
2024	466,557	460,226	101.4%	
Total	7,468,534	7,476,955	99.9%	6,331,403

Table 5.2.1. Catches (tonnes) of Norwegian spring spawning herring by zones for each year 2013-2024. Change from EU to UK EEZ from 2021.

Year	Faroes	Greenland	Iceland	JanMayen	NEAFC	Norway	Russia	Svalbard	UK	Total
2013	118,526	2,221	60,373	16,602	73,626	375,887	15	37,525	1	684776
2014	38,919	11,015	45,418	5,008	31,901	289,665	0	26,961	312	449199
2015	41,893	10,026	39,472	656	20,757	202,007	0	4,735	582	320128
2016	59,456	16,262	48,624	1	46,649	213,293	0	0	411	384697
2017	100,113	9,564	56,577	0	281,186	272,639	0	0	0	720081
2018	64,363	329	46,013	0	164,886	318,676	0	0	100	594367
2019	48,655	0	115,238	0	281,458	328,211	1	0	1	773565
2020	31,780	3	171,390	0	100,912	416,680	0	0	30	720796
2021	60,320	0	164,425	6	30,338	592,911	1	0	97	848099
2022	26,153	1	164,258	29	91,616	545,615	0	13	1	827686
2023	32,366	1	144,382	15	48,386	462,247	1	11	2	687409
2024	5,637	1,375	102,802	31	52,229	304,189	0	3	290	466557

Table 5.2.2. Catches (percentages) of Norwegian spring spawning herring by zones for each year 2013-2024. Change from EU to UK EEZ from 2021.

Year	Faroes	Greenland	Iceland	JanMayen	NEAFC	Norway	Russia	Svalbard	UK
2013	17.3	0.3	8.8	2.4	10.8	54.9	0.0	5.5	0.0
2014	8.7	2.5	10.1	1.1	7.1	64.5	0.0	6.0	0.1
2015	13.1	3.1	12.3	0.2	6.5	63.1	0.0	1.5	0.2
2016	15.5	4.2	12.6	0.0	12.1	55.4	0.0	0.0	0.1
2017	13.9	1.3	7.9	0.0	39.0	37.9	0.0	0.0	0.0
2018	10.8	0.1	7.7	0.0	27.7	53.6	0.0	0.0	0.0
2019	6.3	0.0	14.9	0.0	36.4	42.4	0.0	0.0	0.0
2020	4.4	0.0	23.8	0.0	14.0	57.8	0.0	0.0	0.0
2021	7.1	0.0	19.4	0.0	3.6	69.9	0.0	0.0	0.0
2022	3.2	0.0	19.8	0.0	11.1	65.9	0.0	0.0	0.0
2023	4.7	0.0	21.0	0.0	7.0	67.2	0.0	0.0	0.0
2024	1.2	0.3	22.0	0.0	11.2	65.2	0.0	0.0	0.1

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2013. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	0	1 922	0	2 322	0	4 244
DNK	0	0	0	0	0	17 159	0	0	1	17 160
FRO	97 741	0	0	0	7 297	0	0	0	0	105 038
GRL	2 070	2 221	3	0	7 840	0	0	0	0	12 133
IRL	0	0	0	0	0	3 165	0	0	0	3 165
ISL	18 534	0	60 371	0	11 798	0	0	0	0	90 703
NLD	0	0	0	0	1	2 317	0	3 302	0	5 620
NOR	0	0	0	0	36 549	323 302	0	0	0	359 851
RUS	181	0	0	16 602	10 141	19 681	15	31 901	0	78 521
UK	0	0	0	0	0	8 342	0	0	0	8 342
Total	118 526	2 221	60 373	16 602	73 626	375 887	15	37 525	0.931	684 776

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2014. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	0	384	0	284	0	669
DNK	0	0	0	0	8 893	2 916	0	0	0	11 810
FRO	24 093	0	0	0	2 806	0	0	0	0	26 899
GRL	2 022	11 001	165	0	0	0	0	0	0	13 188
IRL	0	0	0	0	706	0	0	0	0	706
ISL	6 978	14	45 253	0	6 583	0	0	0	0	58 827
NLD	0	0	0	0	72	2 607	0	6 453	312	9 445
NOR	0	0	0	28	7 255	255 847	0	0	0	263 131
RUS	5 826	0	0	4 980	5 586	23 676	0	20 224	0	60 292
UK	0	0	0	0	0	4 233	0	0	0	4 233
Total	38 919	11 015	45 418	5 008	31 901	289 665	0	26 961	312	449 199

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2015. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	1 110	0	0	0	1 547	0	0	0	4	2 660
DNK	649	0	0	0	7 329	14	0	0	0	7 992
FRO	23 053	0	0	0	2 897	0	0	0	180	26 130
GRL	2 058	10 026	231	0	0	124	0	0	135	12 574
IRL	0	0	0	0	999	400	0	0	0	1 400
ISL	3 139	0	39 241	0	246	0	0	0	0	42 626
NLD	2 453	0	0	0	1 977	0	0	0	233	4 663
NOR	0	0	0	0	0	176 176	0	0	0	176 176
RUS	9 433	0	0	656	5 743	25 286	0	4 735	0	45 853
UK	0	0	0	0	19	6	0	0	30	55
Total	41 893	10 026	39 472	656	20 757	202 007	0	4 735	582	320 128

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2016. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	700	0	0	0	1 882	0	0	0	0	2 583
DNK	1 895	0	0	0	219	7 634	0	0	0	9 747
FRO	42 898	0	0	0	1 829	0	0	0	0	44 727
GRL	2 350	16 262	1 495	0	0	0	0	0	0	20 108
IRL	0	0	0	0	1 529	519	0	0	0	2 048
ISL	1 210	0	47 129	0	2 119	0	0	0	0	50 458
NLD	520	0	0	0	1 726	570	0	0	302	3 119
NOR	0	0	0	0	12 341	185 081	0	0	0	197 421
RUS	9 884	0	0	1	24 982	15 589	0	0	0	50 456
UK	0	0	0	0	22	3 900	0	0	109	4 031
Total	59 456	16 262	48 624	1	46 649	213 293	0	0	411	384 697

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2017. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	557	0	0	0	4 607	2	0	0	0	5 166
DNK	0	0	0	0	3 927	13 923	0	0	0	17 850
FRO	57 775	0	0	0	40 388	0	0	0	0	98 163
GRL	2 500	9 564	457	0	48	0	0	0	0	12 570
IRL	1 291	0	0	0	1 179	1 024	0	0	0	3 495
ISL	28 813	0	56 119	0	5 468	0	0	0	0	90 400
NLD	631	0	0	0	5 676	0	0	0	0	6 307
NOR	0	0	0	0	157 794	231 589	0	0	0	389 383
POL	0	0	0	0	0	1	0	0	0	1
RUS	8 514	0	0	0	61 301	21 303	0	0	0	91 118
SWE	0	0	0	0	766	439	0	0	0	1 205
UK	33	0	0	0	31	4 358	0	0	0	4 422
Total	100 113	9 564	56 577	0	281 186	272 639	0	0	0	720 081

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2018. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	1 919	0	0	0	2	1 922
DNK	0	0	0	0	2 094	14 958	0	0	0	17 052
FRO	32 377	0	4 920	0	44 155	510	0	0	0	81 962
GRL	2 460	329	10	0	91	0	0	0	0	2 890
IRL	0	0	297	0	122	2 009	0	0	0	2 428
ISL	19 149	0	40 786	0	23 457	0	0	0	0	83 392
NLD	1 557	0	0	0	2 846	0	0	0	32	4 435
NOR	0	0	0	0	34 849	297 177	0	0	0	332 027
POL	50	0	0	0	932	0	0	0	65	1 047
RUS	8 769	0	0	0	54 421	995	0	0	0	64 185
SWE	0	0	0	0	0	445	0	0	0	445
UK	0	0	0	0	0	2 582	0	0	0	2 582
Total	64 363	329	46 013	0	164 886	318 676	0	0	100	594 367

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2019. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	4 142	0	0	0	0	4 142
DNK	227	0	0	0	11 049	6 413	0	0	0	17 688
FRO	26 664	0	37 685	0	49 591	0	0	0	0	113 940
GRL	1 747	0	0	0	1 677	0	0	0	0	3 424
IRL	0	0	0	0	1 098	1 677	0	0	0	2 775
ISL	11 152	0	77 553	0	19 341	0	0	0	0	108 046
NLD	512	0	0	0	4 647	0	0	0	1	5 160
NOR	0	0	0	0	113 309	317 195	0	0	0	430 504
POL	0	0	0	0	1 327	0	0	0	0	1 327
RUS	8 353	0	0	0	75 278	406	1	0	0	84 039
SWE	0	0	0	0	0	720	0	0	0	720
UK	0	0	0	0	0	1 801	0	0	0	1 801
Total	48 655	0	115 238	0	281 458	328 211	1	0	1	773 565

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2020. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	27	0	0		2 747	0			28	2 801
DNK	0	0	0		2 974	9 885			0	12 860
FRO	21 720	0	77 363		3 947	0			0	103 029
GRL	1 127	3	1		2 419	0			0	3 550
IRL	0	0	0		1 102	1 601			0	2 704
ISL	204	0	94 027		3 655	287			0	98 173
NLD	501	0	0		3 275	3 039			0	6 815
NOR	0	0	0		29 043	380 394			0	409 436
POL	0	0	0		263	1 089			0	1 352
RUS	8 201	0	0		51 121	15 603			0	74 925
SWE	0	0	0		223	2 947			0	3 170
UK	0	0	0		143	1 836			2	1 981
Total	31 780	3	171 390		100 912	416 680			30	720 796

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2021. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	0	3,185	0	0	0	3,186
DNK	0	0	0	0	0	15,854	0	0	0	15,854
FRO	44,821	0	55,779	0	13,682	9	0	0	0	114,290
GRL	6,244	0	1	0	211	0	0	0	0	6,456
IRL	0	0	0	0	0	1,793	0	0	0	1,793
ISL	16	0	108,646	4	5,611	22	0	0	0	114,299
NLD	0	0	0	0	0	7,389	0	0	0	7,389
NOR	0	0	0	0	843	488,790	0	0	0	489,633
POL	0	0	0	0	0	1,144	0	0	97	1,242
RUS	9,240	0	0	1	9,991	73,609	1	0	0	92,842
SWE	0	0	0	0	0	1,116	0	0	0	1,116
Total	60,320	0	164,425	6	30,338	592,911	1	0	97	848,099

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2022. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	0	5,379	0	0	0	5,379
DNK	0	0	0	0	756	14,257	0	0	0	15,014
FRO	16,397	0	60,889	0	44,796	0	0	0	0	122,082
GRL	598	0	0	0	19,989	0	0	0	0	20,587
IRL	0	0	0	0	0	3,209	0	0	0	3,209
ISL	0	0	103,369	0	9,370	0	0	0	0	112,739
NLD	0	0	0	0	1,741	1,873	0	0	0	3,615
NOR	1	1	0	29	2,955	443,281	0	13	1	446,281
POL	0	0	0	0	0	0	0	0	0	0
RUS	9,156	0	0	0	12,008	64,705	0	0	0	85,870
SWE	0	0	0	0	0	3,291	0	0	0	3,291
UK	0	0	0	0	0	9,620	0	0	0	9,620
Total	26,153	1	164,258	29	91,616	545,615	0	13	1	827,686

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2023. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	0	2,109	0	0	0	2,109
DNK	0	0	0	0	8,048	2,550	0	0	0	10,598
FRO	20,070	0	54,289	0	16,012	0	0	0	0	90,371
GRL	3,457	0	0	0	7,167	0	0	0	0	10,623
IRL	0	0	0	0	0	1,016	0	0	0	1,016
ISL	0	0	90,092	0	2,105	0	0	0	0	92,197
NLD	0	0	0	0	0	5,920	0	0	0	5,920
NOR	0	1	0	15	10	389,453	0	11	2	389,491
POL	0	0	0	0	1,331	0	0	0	0	1,331
RUS	8,840	0	0	0	13,713	51,592	1	0	0	74,145
SWE	0	0	0	0	0	781	0	0	0	781
UK	0	0	0	0	0	8,827	0	0	0	8,827
Total	32,366	1	144,382	15	48,386	462,247	1	11	2	687,409

Table 5.2.3. Catches (tonnes) of Norwegian spring spawning herring by zones in 2024. Change from EU to UK EEZ from 2021.

Country \ EEZ	Faroes	Greenland	Iceland	Jan Mayen	NEAFC	Norway	Russia	Svalbard	UK	Total_catch
DEU	0	0	0	0	1,677	2,121	0	0	2	3,799
DNK	0	0	0	0	2,660	4,508	0	0	0	7,168
FRO	79	0	38,149	0	20,348	8	0	0	0	58,584
GRL	0	1,375	2,218	0	6,880	0	0	0	0	10,473
IRL	0	0	0	0	0	2,141	0	0	0	2,141
ISL	0	0	62,435	0	532	0	0	0	0	62,966
NLD	0	0	0	0	0	3,062	0	0	0	3,062
NOR	0	0	0	31	656	258,045	0	3	285	259,021
POL	0	0	0	0	0	0	0	0	3	3
RUS	5,559	0	0	0	19,476	24,755	0	0	0	49,790
SWE	0	0	0	0	0	3,706	0	0	0	3,706
UK	0	0	0	0	0	5,843	0	0	0	5,843
Total	5,637	1,375	102,802	31	52,229	304,189	0	3	290	466,557

9. Figures




Figure 2.1.1. ICES ecoregions and statistical areas (Divisions), where Norwegian spring spawning herring is distributed. Also, adjacent Divisions are shown.




Figure 2.1.1.1. Spawning area of Norwegian Spring Spawning Herring in February 2020. Data from Norwegian Acoustic survey on the spawning grounds.




Figure 2.1.1.2. Migration scheme of the 1983 year-class as juveniles in the Barents Sea (from Røttingen 1990).




Figure 2.2.1. Current migration pattern of the adult part of Norwegian spring-spawning herring (NSSH) and interactions with other surrounding stocks, i.e. Icelandic summer-spawning herring (ISSH), Faroese autumn-spawning herring (FASH), and Norwegian autumn-spawning herring (NASH) (from Pampoulie et al. 2015).




Figure 4.1.1. Distribution of herring from the acoustic survey on the spawning grounds in February 2015-2025 (continues on next page).




Figure 4.1.1. continues.




Figure 4.1.1. continues.




Figure 4.2.1. Distribution of herring larvae in April 2014-2016.




Figure 4.3.1. Distribution of juvenile NSSH in Barents Sea from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2014-2024. Note that for all years except 2016 and 2024, the map covers also the distribution of adult fish in the Nordic Seas. There was no coverage in the Barents Sea in 2020, 2022, 2023 and 2025. The coverage in 2024 was reduced and only the eastern part of the Barents Sea was covered.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2014.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2015.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2016.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2017.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2018.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2019.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2020.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2021.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2022.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2023.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2024.




Figure 4.4.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem survey in the Nordic Seas (IESNS) in May/June 2025.





Figure 4.5.1. Distribution of 0-group NSSH from the ecosystem survey in the Barents Sea in autumn 2014-2023. No maps for 2024 and 2025.

Coastal States WG Herring 2025






Figure 4.5.2. Distribution of juvenile NSSH from the ecosystem survey in the Barents Sea in autumn 2014-2024. No map for 2025 since the survey is not completed.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2014.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2015.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2016.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2017.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2018.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2019.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2020.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2021.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2022.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2023.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2024.




Figure 4.6.1. Distribution of adult NSSH in the Nordic Seas from the International ecosystem summer survey in the Nordic Seas (IESSNS) in July 2025.




Figure 5.1.1. Relative catch by EEZs in the period 1995-2024 (left axis) and total catch mill. tonnes (right axis). *Change from EU to UK EEZ from 2021.




Figure 5.3.1. Relative catch per month for the years 1995-2024. The coloured bold lines are averages over different periods (yellow 1995-1997, green 1998-2005, red 2006-2012, black 2013-2023, and light blue 2024).

Annexes

Annex 1 – Survey and catch tables from the 2014 report

Please refer to the 2024 report for survey and catch tables 1995-2013 (Anon. 2024).

Annex 2. Distribution of NSSH fishing per month and year




Figure A2.1. Total catches combined for years 2013-2024 by month.




Figure A2.2. Total catches 1993-2024 in January.




Figure A2.2. Total catches 1993-2024 in February.




Figure A2.2. Total catches 1993-2024 in March.




Figure A2.2. Total catches 1993-2024 in April.




Figure A2.2. Total catches 1993-2024 in May.




Figure A2.2. Total catches 1993-2024 in June.




Figure A2.2. Total catches 1993-2024 in July.




Figure A2.2. Total catches 1993-2024 in August.




Figure A2.2. Total catches 1993-2024 in September.




Figure A2.2. Total catches 1993-2024 in October.




Figure A2.2. Total catches 1993-2024 in November.




Figure A2.2. Total catches 1993-2024 in December.

Annex 3. Data call and formats

As stated in the agreed record from London 2024 on management of Norwegian spring-spawning herring, the coastal States agreed to update the 'Report of the Coastal states Working Group on the distribution of Norwegian Spring Spawning Herring in the Northeast Atlantic and the Barents Sea' in 2025, chaired by Faroes.

The survey data will be accessible from the survey groups. However, updating the catch data requires your attention now, and is the reason for this data call.

We therefore request you all to provide information on your respective catches/by-catches of NSSH by ICES statistical squares and months, and by exclusive economic zones (EEZs) and international waters, for 2024 (see details in the attached document). We kindly ask you to provide this data by 1 June 2025.

This information should be submitted to Jan Arge Jacobsen (janarge@hav.fo), who will be responsible for compiling the information and make it available for the participants.

The WG will aim for an online meeting in early autumn to finish the updated report. If anyone wants the WG to meet before then to clarify issues, please be in contact. Please inform Jan Arge of additional participant you think should be added.

Content of attached document:

Catch data should be submitted in a text file with 7 columns:
year, species, country, ices_rect, month, catch, zone

It is important that:

- The columns are comma-separated
- The parameters are put in the file without any "
- The file is saved as .txt or .csv
- Only upper case letters are used
 - Countries are indicated with the three letterscode (alpha-3 code, https://www.nationsonline.org/oneworld/country_code_list.htm). **See separate designations for the United Kingdom at the bottom of the Table below.**
- Rectangle names are without spaces or hyphens
- Month is given in numbers: 1,2, ..., 12
- Landings/catches are given in tonnes (with three decimal places and use *point* as the decimal separator (not *coma*) e.g. 15000.123)
- Economic zones are given as three letter codes (see examples below)

Example of submission file. (The first line is the header line):

```
year,species,country,ices_rect,month,catch, zone
2021,HER,DEU,55E8,3,99.000,IRL
2021,HER,DEU,55E8,4,4.210,IRL
2021,HER,DEU,55E9,4,54.321,IRL
```

(catch numbers given in the example do not correspond to true values)

From EEZ / MarineRegions:


```
Territory1                ISO_Ter1
-----
```

Coastal States WG Herring 2025

International Barents Sea	INB
International North	INN
International West	INW
Belgium	BEL
Germany	DEU
Denmark	DNK
Spain	ESP
Alhucemas Islands	ESP
Perejil Island	ESP
Ceuta	ESP
Peñón de Vélez de la Gomera	ESP
Chafarinas Islands	ESP
France	FRA
Faeroe	FRO
United Kingdom*	GBR
Guernsey	GGY
Gibraltar	GIB
Greenland	GRL
Ireland	IRL
Iceland	ISL
Jersey	JEY
Netherlands	NLD
Norway	NOR
Poland	POL
Azores	PRT
Madeira	PRT
Portugal	PRT
Russia	RUS
Svalbard	SVA
Jan Mayen	SJM
Sweden	SWE

*Separate regions in the United Kingdom

UK/Scotland	UKS
UK/England	UKE
UK/Wales	UKW
UK/Northern Ireland	UKN



Annex 4. List of participants

Åge Høines, Norway

Alexander I. Krysov, Russia

Erling Kåre Stenevik, Norway

Guðmundur J. Óskarsson, Iceland (chair)

Sigurvin Bjarnason, Iceland

Jan Arge Jacobsen, Faroe Island

Eydna í Homrum, Faroe Island

Søren Post, Greenland

Richard Nash, UK

Christine Rockmann, EU Commission